

All investigation showed no evidence of tumor in ovaries and at any other primary site then the patient finally diagnosed as having Granulosa cell tumor involving only omentum post op stage III C. Then patient was given six courses of chemotherapy with Inj Paclitaxel and Inj Carboplatin three weekly. Now patient is on regular follow up and disease free.

**Conclusion:** Extra ovarian adult granulosa cell tumor of omentum is rare tumor. Multimodal treatment approaches including surgery, multi-agent chemotherapy may provide a survival benefit for patients.

## Miscellaneous: Ovary Abstract

### Synchronous malignancy of ovary and cervix

**Vani Malhotra, Smriti Nanda, Meenakshi Chauhan, Vandana Bhuria**

**Background:** Synchronous primary malignancies of the female genital tract constitute 1.7% of all genital malignancies.

**Case:** A 45-year-old para 5 woman presented with loss of appetite and abdominal distention. Provisional diagnosis of ovarian malignancy was made. Final histopathology of the specimen revealed ovarian papillary serous cystadenocarcinoma with cervical leiomyosarcoma. She received chemotherapy.

**Results:** Patient is on regular follow-up.

**Conclusion:** The coexistence of primary neoplasms in the ovary and cervix are rare. A normal appearing organ may have a hidden malignancy. So, every surgical specimen should be subjected to detailed histopathological examination. Also, the possibility of synchronous malignancy elsewhere in body should be kept in mind while working on a genital malignancy.

## Miscellaneous: Poster Abstract

### Radical excision of a massive vulvo: Vaginal mass

**J. Meena, A. Parthasarathy, R. Vatsa, N. Singh, S. Kumar, K. K. Roy, S. Singhal**

**Department of Obstetrics and Gynaecology, AIIMS, New Delhi, India**

**Background:** Vulvo-vaginal masses has a varied presentation and causes. The most common differential diagnosis are Condylomata acuminata, Vulvar abscess, Vulvar/vaginal cysts, Vulval carcinoma, Vulval lipoma, Angiomyofibroblastoma and Aggressive Angiomyxoma. Surgical excision of the mass is the main method of treatment and the outcome differs with the histological diagnosis. We present a video of excision of a massive Vulvo vaginal mass in toto.

**Case:** A 45 year old P3 L3 female, presented with complaint of mass in perineal area & discharge per vaginum for 2 years. The mass was growing progressively and reached the present size. On examination there was a 9 X 8 cm irregular firm to cystic mass, arising from posterior wall of vagina and protruding out of introitus with bossellated surface. The mass also extended into right ischio-rectal fossa, 10 X 10 cm mass with cystic, smooth surface that was irreducible with no cough impulse. CECT abdomen and pelvis revealed a well-defined 12 X 10 X 8 cm mass in right perineum arising from right lower lateral vaginal wall with ischio-rectal fossa extension. There was no extension into cervix, bladder or rectum. Biopsy taken from the mass was inconclusive. A wide local excision was done under general anesthesia wherein an ischio-rectal and vaginal mass of size 30 X 10 cm with irregular margin was excised in toto. Histopathology was suggestive of Aggressive Angiomyxoma. The patient is under follow up.

**Discussion:** Aggressive Angiomyxoma is a rare slow growing locally invasive mesenchymal tumor and has a substantial potential for recurrence. It is often misdiagnosed. Pre-operative diagnosis is difficult due to rarity of this entity and absence of diagnostic features, but it should be considered in case of masses in genital, perianal and pelvic region in a woman of reproductive age. Radical surgical excision is the first line of management. A long term follow up of the case is necessary and MRI is preferred method for detecting recurrences.

## Miscellaneous: Poster Abstract

### Distant pedicled musculocutaneous/fasciocutaneous flaps; a novel approach for reconstruction of large vulvar defects

**Kripa Shanker Mishra**

## RGCIIRC, New Delhi, India

**Introduction:** Postablative reconstruction of vulvar defects is a difficult challenge. Local flaps carry a high incidence of delayed wound healing as local flaps may redistribute but not eliminate local wound tension. Repair of defect with distant pedicled flaps may avert local complications by minimising tension to the skin and increasing the initial biomechanical strength of wound. The aim of this study was to determine the clinical outcome of distant musculocutaneous & fasciocutaneous flaps used for postablative reconstruction of large vulvar defects.

**Methods:** Between January 2015 to December 2015 total three patients underwent vulvectomy and immediate reconstruction with distant pedicled flaps for vulvar carcinoma. Postoperative complications were recorded and clinical outcomes were evaluated.

**Results:** Two of the three flaps healed primarily. One flap was complicated by minor wound dehiscence, which healed with conservative treatment. Hospital stay and clinical course was shorter in comparison to local flaps and split skin graft.

**Conclusions:** Distant musculocutaneous and fasciocutaneous flaps provide excellent design flexibility and can be tailored to reconstruct postablative vulvar defects with good outcomes and minimal morbidity.

## Miscellaneous: Poster Abstract

### To find the prevalence of female genital tract malignancies in a tertiary care hospital

**Dilpreet K. Pandher**

**Department of Obstetrics and Gynaecology, Government Medical College and Hospital, Sector 32, Chandigarh, India**

Genital tract and breast are two most common sites of malignancy in females. Out of the genital tract malignancies, carcinoma cervix is so far found to be the commonest followed by ovary and endometrium. In developed countries, carcinoma cervix incidence is comparatively quite low due to good regular screening of females. One year review of patients was done, who underwent definitive/debulking surgery for a diagnosed malignant pathology of the genital tract, in obstetrics and gynaecology department of Govt medical College and Hospital, Chandigarh. Total 62 patients were operated, most common indication was carcinoma ovary, followed by endometrial cancer, cancer cervix and gestational trophoblastic neoplasia. 166 patients underwent biopsies for suspicious symptoms or the abnormal findings on examination and the patients with final malignancy report were either operated as described above and the inoperable cases were referred to oncology department for further management.

## Miscellaneous: Poster Abstract

### Gestational trophoblastic neoplasia: Retrospective analysis of clinical profile, treatment pattern and outcome

**Paramjeet Kaur, Ashok K. Chauhan, Anil Khurana, Yashpal Verma, Nupur Bansal**

**Department of Radiotherapy, Pt. BDS PGIMS, Rohtak, Haryana, India**

**Background:** Gestational trophoblastic disease is a spectrum of cellular proliferation arising from the placental villous trophoblast. Gestational trophoblastic neoplasia (GTN) is a collective term for GTD that invade locally or metastasize. GTD includes hydatidiform mole (complete and partial) and GTN include invasive mole, choriocarcinoma, placental site trophoblastic tumor and epithelioid trophoblastic tumor.

**Aim:** To evaluate clinicopathological profile, treatment pattern and clinical outcome in patients with gestational trophoblastic neoplasia (GTN).

**Materials and Methods:** Twelve cases of gestational trophoblastic neoplasia treated between 2012 to November 2015 in deptt of Radiotherapy – II, PGIMS, Rohtak were evaluated in this retrospective study. Data was analyzed on the basis of age, histopathology, stage, type of treatment received and treatment related toxicities. Disease free survival was estimated.

**Results:** Out of 12 women 7 (58 %) had hydatidiform mole, 4 (33%) invasive mole and 01 (8%) had choriocarcinoma. All the cases were given chemotherapy. Two patients had low risk disease. Among high risk group