

Cervix: Poster Abstract

Chemoradiation for the management of locally advanced carcinoma uterine cervix: Comparative evaluation of concomitant weekly versus three weekly cisplatin

Sulbha Mittal

Department of Radiation Oncology, Pt. B.D. Sharma PGIMS, Rohtak, Haryana, India

Aims and Objectives: To determine and evaluate the difference/s, in terms of tumor control and side effects, between weekly and three weekly cisplatin concomitant with external beam radiotherapy for locally advanced carcinoma of cervix.

Materials and Methods: The study was conducted in Radiotherapy Department, University of Health Sciences, Rohtak (India), on sixty previously untreated, histopathologically proven patients of locally advanced carcinoma of uterine cervix. The patients were treated with External Beam Radiotherapy (EBRT) 50 Gy/25 fractions over 5 weeks and concomitant cisplatin, followed by intra-cavitary HDR brachytherapy (ICBT) 700 cGy to point A; three times, once in a week. The patients were assigned randomly either of two groups of 30 patients each. In Group I (Study Group) the patients received three weekly cisplatin 75 mg/m² for 2 cycles while in Group II (Control Group) the patients received weekly cisplatin 40 mg/m² for 5 cycles. Evaluation of response and toxicity was done weekly during treatment and monthly thereafter up to six months. The data thus obtained was assessed and analysed using LaMorte statistical tool. The study was approved by Ethical Committee of the institute and quality was periodically monitored by senior consultant and guide.

Results: Stage wise disease response in study and control respectively at the end of treatment was as follows: Stage IIA - CR (80% vs 100%), PR (20% vs 0%); Stage IIB - CR (80% vs 76.47%), PR (20% vs 23.53%); Stage IIIA - CR (60% vs 100%), PR (40% vs 0%); Stage IIIB - CR (60% vs 60%), PR (40% vs 20%), NR(0% vs 20%). Stage wise disease status at the end of sixth month follow up was as follows: Stage IIA - NED (80% vs 100%), RD (20% vs 0%); Stage IIB - NED (80% vs 76.67%), RD (20% vs 23.53%); Stage IIIA - NED (60% vs 100%), RD (40% vs 0%); Stage IIIB - NED (60% vs 60%), RD (40% vs 40%). Tumor response was not significantly different in the two groups with respect to age distribution, rural/ urban distribution, histopathological distribution and treatment interruption. Maximum level of hematological toxicity (WHO criteria) observed in study and control group respectively at the end of treatment was as follows: Anaemia; Grade II - 4 (13.33%) in both the groups, leukopenia; Grade II - 1 (3.33%) vs 0 (0%). The worst acute skin reactions observed by the end of treatment in Group I and II respectively were Grade II - 2 (6.67%) vs 0 (0%). The worst acute mucosal reactions were Grade II - 5 (16.66%) vs 0 (0%). Upper gastrointestinal toxicity (Grade II & III) was 16.7% versus 13.3% respectively. Lower gastrointestinal toxicity (Grade II & III) was 30.0% versus 36.7%. No significant weight loss was observed in either of the groups. Though, all the patients completed the intended treatment, treatment interruption for more than a week was observed in 10 (33.33%) vs 8 (26.67%) patients respectively, due to acute toxicities.

Conclusion: Three weekly cisplatin, concomitant with radiation seems to be the potential, effective and acceptable alternate as standard of treatment for locally advanced carcinoma cervix; especially for increased work load and limited resource setups.

Cervix: Poster Abstract

Comparison of Keyes punch biopsy instrument with cervical punch biopsy forceps for diagnosing cervical lesions

Natasha Tyagi, Amita Suneja, Kiran Mishra¹, Sandhya Jain, Neelam B. Vaid, Kiran Guleria

Departments of Obstetrics and Gynaecology and ¹Pathology, UCMS and GTB Hospital, New Delhi, India

Aims: To assess the feasibility and efficacy of Keyes punch biopsy instrument (KP) in diagnosing cervical lesions and compare it with cervical punch biopsy forceps (CP).

Methods: 75 women having adequate colposcopy with abnormal transformation zone were included and paired colposcopic directed biopsies were taken using KP followed by CP from the same target area. The outcome parameters were compared using paired t-test, Wilcoxon signed rank test and McNemar test.

Results: It was feasible in all cases to take cervical biopsy with KP and CP. Volume of gross specimen obtained by KP was less than CP (0.076±0.097 vs 0.101±0.156 cm³, p=0.061), however on microscopic examination, mean length and depth of tissue in KP was greater than CP by 0.06 mm (p=0.810) and 0.14 mm (p=0.634) respectively. There was an exact agreement with final surgical specimen in 42% of cases in both forceps. Agreement within 1 degree was found in 25% of cases with KP and in 17% of cases with CP. Both the forceps equally missed microinvasive lesions but KP was inferior to CP for invasive cancer.

Conclusion: KP is almost at par with CP for diagnosing preinvasive cervical lesions and is a useful adjunct to the existing armamentarium of biopsy forceps.

Cervix: Oral Abstract

The impact of tumour regression in locally advanced carcinoma cervix during external beam radiotherapy and the need for adaptive planning

Aim: To study the impact of tumour regression occurring during IMRT for locally advanced carcinoma cervix and study dose distribution to target volume and OARs and hence the need for any replanning.

Materials and Methods: 40 patients undergoing IM-IGRT and weekly chemotherapy were included in the study. After 36 Gy, a second planning CT-scan was done and target volume and OARs were recontoured. First plan (non-adaptive) was compared with second plan (adaptive plan) to evaluate whether it would still offer sufficient target coverage to the CTV and spare the OARs after having delivered 36 Gy. Finally new plan was created based on CT-images to investigate whether creating a new treatment plan would optimize target coverage and critical organ sparing. To measure the response of the primary tumour and pathologic nodes to EBRT, the differences in the volumes of the primary GTV and nodal GTV between the pretreatment and intratreatment CT images was calculated. Second intratreatment IMRT plans was generated, using the delineations of the intratreatment CT images. The first IMRT plan (based on the first CT-scan or non adaptive plan) was compared with second IMRT plan (based on the second CT-scan or adaptive plan).

Results: 35% patients had regression in GTV in the range of 4.1% to 5%, 20% in the range of 1.1%-2%, 15% in the range of 2.1%-3% and 20% in the range of 6%-15%. There was significant mean decrease in GTV of 4.63 cc (p=0.000). There was a significant decrease in CTV on repeat CT done after 36 Gy by 23.31 cc (p=0.000) and in PTV by 23.31 cc (p=0.000). There was a statistically significant increase in CTV D98, CTV D95, CTV D50 and CTV D2 in repeat planning CT done after 36 Gy. There was no significant alteration in OARs doses.

Conclusion: Despite tumour regression and increased target coverage in locally advanced carcinoma cervix after a delivery of 36 Gy there was no sparing of OARs. Primary advantage of adaptive RT seems to be in greater target coverage with non-significant normal tissue sparing.

Ovary: Oral Abstract

A prospective study evaluating preoperative (clinical, imaging) and intraoperative predictors of suboptimal debulking in advanced epithelial ovarian cancer

Anupama Rajanbabu, Kiran Bagul, Venkatesan

Department of Surgical Oncology, Amrita Institute

Introduction: In advanced epithelial ovarian cancer, there is a survival benefit for patients who achieve optimal cytoreduction. Suboptimally cytoreduced patients are at risk of the increased morbidity of a surgery without associated survival benefit. Predicting which patients can undergo optimal cytoreduction represents a critical decision-making point. Present study analyses the predictors, pre operative (clinical and radiologic) and intraoperative of suboptimal debulking.

Methods: This was a prospective observational study conducted at Amrita Institute of Medical Sciences from March 2013 to May 2015. All the patients with clinically (physical examination, laboratory and imaging results) diagnosed Stage IIIc epithelial ovarian, fallopian tube, or primary peritoneal carcinoma (PPC) who were planned for primary debulking surgery were included. The demographic data and details of tumor markers, radiological investigations including CT scan, intra operative findings and histopathologic details were collected. Statistical analysis was done using SPSS v20.0.